


Seventh Semiannual Progress Report  
September, 1974 - February, 1975  
for  
National Aeronautics and Space Administration  
Grant NGL 15-005-140  
entitled

Origin of Chondrites from Relationships  
Between and Thermal Retentivity of  
Key Trace Elements

awarded to  
Purdue Research Foundation  
Purdue University  
West Lafayette, Indiana 47907  
prepared by  
Michael E. Lipschutz, Principal Investigator  
Professor of Chemistry and Geosciences  
Purdue University  
West Lafayette, Indiana 47907



Date of Report: February 28, 1975

(NASA-CR-142620) : ORIGIN OF CHONDRITES FROM  
RELATIONSHIPS BETWEEN AND THERMAL  
RETENTIVITY OF KEY TRACE ELEMENTS  
Semiannual Progress Report, Sep. 1974 - Feb.  
1975 (Purdue Univ.) : 8 p.

N75-73000

Unclas  
10211

REPRODUCED BY  
NATIONAL TECHNICAL  
INFORMATION SERVICE  
U.S. DEPARTMENT OF COMMERCE  
SPRINGFIELD, VA 22161

00/98

## I. General Remarks

Since submission of our Sixth Semiannual Progress Report (March - August, 1974) the following papers have been published, are in press or have been submitted for publication:

- (1) "Trace Elements in Primitive Meteorites-V. Abundance Patterns of Thirteen Trace Elements and Interelement Relationships in Enstatite Chondrites," C. M. Binz, R. K. Kurimoto and M. E. Lipschutz, Geochimica et Cosmochimica Acta, 38, 1579-1606 (1974).
- (2) "R-Mode Factor Analysis on Enstatite Chondrite Analyses," D. M. Shaw, Geochimica et Cosmochimica Acta, 38, 1607-1613 (1974).
- (3) "Thermal Metamorphism of Primitive Meteorites-I. Variation of Six Trace Elements in Allende Carbonaceous Chondrite Heated at 400-1000°C," M. Ikramuddin and M. E. Lipschutz, Geochimica et Cosmochimica Acta, 39, 363-375 (1975).
- (4) "Allende Meteorite : Effect of Thermal Metamorphic Conditions on Mineralogy and Trace Element Retention," M. Ikramuddin, M. E. Lipschutz and W. R. Van Schmus, Nature (London), in press (1975).
- (5) "Trace Elements in Primitive Meteorites-VI. Abundance Patterns of Thirteen Trace Elements and Interelement Relationships in Unequilibrated Ordinary Chondrites," C. M. Binz, M. Ikramuddin, P. Rey and M. E. Lipschutz, Geochimica et Cosmochimica Acta, submitted (1975).
- (6) "Contents of Eleven Trace Elements in Ureilite Achondrites," C. M. Binz, M. Ikramuddin and M. E. Lipschutz, Geochimica et Cosmochimica Acta, submitted (1975).

As noted in the Sixth Semiannual Progress Report paper (2) above does not acknowledge NASA support but it is based upon research (i.e. paper (1)) supp-

orted by this grant.

## II. Trace Element Studies

As discussed in the Fourth (March-August, 1973) Fifth (September, 1973-February, 1974) and Sixth Semiannual Progress Reports, our investigations are proceeding along two major avenues: studying effects of extended heating (i.e. simulated metamorphism) of primitive meteorites; establishing key trace element contents in primitive chondrites and meteorites suspected of being derived directly from them.

### A. Laboratory Simulation of Thermal Metamorphic Effects

Since this avenue represents a new approach to trace element studies we continue to emphasize this strongly. In the first paper reporting such experiments, paper (3) cited in part I, we report that determination by neutron activation of 6 trace elements retained in Allende (C3) samples heated at 400-1000°C for 1 week in a low-pressure (initially  $\sim 10^{-5}$  atm)  $H_2$  atmosphere reveals loss of small proportions of Ga and Se and large proportions of Bi, In and Tl - Co being unaffected. The retentivity patterns for the 5 mobile elements differ and in no way duplicate the step-function assumed previously. In contrast to these trace elements sulfur is initially present in discrete mineral(s) and visually it appears to be released over a narrow temperature range. Elements are lost more easily from powder than from chips but the difference is  $\leq 35\%$ . Above 600°C, the process of loss appears due to process(es) with apparent activation energies of 2 kcal/mole (Bi, Tl), 4 kcal/mole (Se) and 22 kcal/mole (In). Loss of Bi and Tl below 600°C involves higher apparent activation energies. Two-element correlation diagrams involving Bi, In and Tl are consistent with the idea that trends among highly-volatile elements in enstatite chondrites arise from metamorphism.

In the second of our studies, paper (4) cited in part I, we report that heating Allende carbonaceous chondrite in a low-pressure environment causes mineralogic alteration at 700-1000°C but not at  $T \leq 600^\circ\text{C}$ . Samples heated for 29 days at 500°C lose Bi, In and Tl more effectively than those similarly heated for 7 days. At 1000°C with  $\sim 10^{-5}$  atm. initial pressure of  $\text{O}_2$ ,  $\text{H}_2$  or He these elements, Ga and Se are comparatively more completely lost. These results indicate the necessity for additional experiments in order to study elemental loss patterns in primitive objects.

Such studies are now in progress. We have completed determination of 10 trace elements (Ag, Bi, Co, Cs, Ga, In, Se, Te, Tl and Zn) in samples at 100° increments over the 400-1000°C range, including duplicates at 800° and 900°C with a molecular sieve in the cold trap to study internal atmospheric effects. We have also analyzed these data statistically (by factor analysis and to establish the pattern of significant interelement relationships); all results are consistent with the idea that enstatite chondrites reflect metamorphic effects. We are currently preparing a manuscript describing these results.

We have completed trace-element measurements of BCR-1 powder heated at 100°C increments over the 500-1000°C temperature span. This study is aimed at comparing trace element retentivity in a sample formed at high temperatures with retentivities in low-temperature accumulates like primitive meteorites. The elements studied include Ag, Bi, Cd, Co, Cs, Ga, In, Se, Tl and Zn (Te was below our detection limit). While most elements yielded coherent trends, Ag, Cd and Se were erratic; we now suspect the cause for this, and plan a series of experiments using BCR-1 chips similarly heated.

We have completed heating samples (at 100°C increments over the 400-1000°C range) of an unequilibrated ordinary chondrite (UOC) and will shortly begin

# DOCUMENT REQUEST

NASA SCIENTIFIC AND TECHNICAL INFORMATION FACILITY

OPERATED BY INFORMATICS INFORMATION SYSTEMS COMPANY

POST OFFICE BOX 8757 BALTIMORE WASHINGTON INTERNATIONAL AIRPORT, MARYLAND 21240 TELEPHONE (301) 796 5300

## DOCUMENT REQUESTED

A NASA ACCESSION NUMBER  N75 73000	B NACA/NASA REPORT NUMBER  OR
C (PLEASE DO NOT WRITE IN THIS SPACE)	
D COPY TYPE REQUESTED: case file <input type="checkbox"/> MICROFICHE <input checked="" type="checkbox"/> FULL SIZE	

## REQUESTER IDENTIFICATION

E REQUESTER'S FACILITY IDENT NO  2523	F REQUESTER'S CONTRACT NO.
G AUTHORIZED SIGNATURE AND DATE  Ellen Davis May 29, 1981	

## OTHER BIBLIOGRAPHIC INFORMATION (ESSENTIAL IF ITEMS A AND B ARE UNKNOWN)

H. DOCUMENT TITLE	
I DATE OF REPORT 04 JAN 1981	
K CORPORATE SOURCE	L CORPORATE REPORT NO.
	M.CONTRACT NO

N. MAILING LABEL (must be imprinted on all copies. include zip code)

Ellen Davis

NTIS

NOTE: For prompt service, please follow instructions on back of last copy

1. SHIPPING COPY

## RESPONSE TO DOCUMENT REQUEST

(See item checked below for the specific reply to your request)

### THE DOCUMENT YOU REQUESTED

#### MAY BE OBTAINED FROM

- ☐ (1) Superintendent of Documents, U S G P O , Washington D C 20401.  
☐ (2) National Technical Information Service, Springfield, Va. 22150  
☐ (3) Defense Documentation Center, Cameron Station, Alexandria, Va 22314  
☐ (4) \_\_\_\_\_

#### IS OUT OF STOCK AND NOT REPRODUCIBLE BECAUSE

- ☐ (5) Copyrighted ☐ (8) Not suitable for reproduction  
☐ (6) Journal Article ☐ (9) Source prohibits reproduction  
☐ (7) Purchase item contact source

#### HAS DISTRIBUTION LIMITATIONS WHICH PREVENT US FROM SATISFYING YOUR REQUEST

Available from the Facility to

- ☐ (10) NASA only. ☐ (12) U S Government Agencies only  
☐ (11) NASA and its contractors only ☐ (13) U S Government Agencies and Contractors only  
☐ (14) Classified. our records do not indicate adequate clearance contact your cognizant contracting agency  
☐ (15) Classified document in Category \_\_\_\_\_ our records do not indicate that your organization has been certified access to that category  
☐ (16) Non-NASA document and therefore available from the Facility only to NASA and its contractors our records do not indicate that you are registered with us as a NASA contractor.  
☐ (17) Source controls and monitors all distribution

#### IS NOT AVAILABLE FOR THE FOLLOWING ADMINISTRATIVE REASON

- ☐ (18) Not available outside U S  
☐ (19) Requires approval of another Government agency for release (Serv. Rept). this approval is being sought; you will be notified.  
☐ (20) Approval sought in #19 has been denied  
☐ (21) Contains proprietary information, requiring approval of responsible NASA Office for release (Spec Rel) this approval is being sought, you will be notified  
☐ (22) Approval sought in #21 has been denied  
☐ (23) Obsolete; withdrawn from circulation  
☐ (24) Out of subject scope, not retained in Facility's collection  
☐ (25) Out of print, not to be reprinted or reproduced  
☐ (26) Repeated attempts to obtain have been unsuccessful

#### IS NOT YET AVAILABLE

Request again when announced in STAR or CSTAR journals

- ☐ (27) Availability is under review ☐ (29) Not yet published  
☐ (28) Review Copy or Advance Copy stage of publication  
☐ (30) Out of stock being reprinted will be forwarded  
☐ (31) Not in Facility's collection action has been taken to obtain copies, you will be notified

#### IS INADEQUATELY IDENTIFIED

- ☐ (32) Please furnish correct NACA/NASA accession number or report number or additional bibliographic information  
☐ (33) Accession number or report number cited is not valid, check reference

#### IS NOT AVAILABLE IN COPY TYPE REQUESTED

- ☐ (34) Available in microfiche only, a microfiche is enclosed  
☐ (35) Available in printed copy only

#### IS NOT AVAILABLE IN MULTIPLE COPIES

- ☐ (36) Enclosed is one photocopy and one microfiche, the microfiche may be utilized as a reproducible master

analyzing these to extend our studies to this third major sort of primitive chondrite.

As in our Allende study, in addition to trace elements, we are continuing to determine the mineralogic and petrologic characteristics of heated samples (Abee, BCR-1, etc.) in cooperation with Professor W. R. Van Schmus of the University of Kansas. We are continuing to study the nature of the organic compounds generated during the heating of meteoritic and geologic samples in cooperation with Professor C. Ponnamperna of the University of Maryland.

We have assembled a system that will enable us to extend the studies above to temperatures in the 1000-1400°C range but, because of the pressure of other studies, we have not yet conducted a systematic study of any sample type. We plan to attend to this in the near future.

#### B. Key Trace Elements in Primitive Meteorites

We have also made substantial progress in this area during the period covered by this report. In paper (1), cited in part I, we describe our determinations of As, Au, Bi, Cd, Co, Cu, Ga, In, Sb, Se, Te, Tl and Zn in 11 samples representing 9 chondrites of grades E4-6. These chondrites exhibit systematic intra- and inter-grade differences particularly for highly-variable elements, the differences being  $E4 \geq E3 > E6 \geq E5$ . The abundance pattern for these 13 and an additional 16 elements in E3-6 chondrites differs from those of other primitive meteorites - the carbonaceous and unequilibrated ordinary chondrites. A search for statistically-significant interelement relationships among the 13 elements (for grades E4-6) reveal that 40 element-pairs are linearly and/or exponentially correlated. Similar consideration of data for 37 elements in 12 chondrites (grades E3-6) reveals that 191 element-pairs exhibit such relationships, 170 involving linear and/or exponential correlations,

the remainder involving anti-correlations. The patterns depicting these relationships - i.e. the correlation profiles - and elemental abundance patterns, factor analysis and two-element correlation diagrams are consistent with all enstatite chondrites representing a single evolutionary sequence. The primary process responsible for the chemical trends of these chondrites involved thermal fractionation accompanied by geochemical fractionation of sulfide-plus-metal from silicate, probably during condensation and accretion of solid material from the solar nebula. Chalcophile elements may have been fractionated during condensation or, after accretion, during thermal metamorphism in the parent body. Indeed as described in paper (3) of Part I, the trends of Bi, In and Tl in "as-received" enstatite chondrites are consistent with those in heated Allende samples. No genetic model proposed thus far accounts for the detailed chemical trends although the constrained equilibrium theory and two-component condensation theories qualitatively seem most satisfactory. The correlation profiles of enstatite, carbonaceous and unequilibrated ordinary chondrites are distinctly different pointing to major differences in the formation conditions of these different sorts of primitive meteorites.

As mentioned, our data were supplied to D. M. Shaw and were used by him for factor analysis. These results were published by him (cf. paper (2) in part I) and appeared simultaneously with our study of the enstatite chondrites.

Paper (5) cited in Part I describes our determinations of As, Au, Bi, Cd, Co, Cu, Ga, In, Sb, Se, Te, Tl and Zn in 13 different unequilibrated ordinary chondrites (UOC), i.e. those having chemically-inhomogeneous silicates. This study, together with prior data for key trace elements, completes coverage of this important group of 23 primitive chondrites. Five elements are quite variable in UOC (As and Cd - 10x, In - 15x, Bi - 300x and Tl - 1000x), the



others varying only by 2-4x. Three highly-depleted elements - Bi, In and Tl - are richer by 5-35x in unequilibrated chondrites than in their equilibrated congeners. All 3 elements vary directly in characteristic fashion with disequilibrium parameters for olivine and pyroxene in UOC and generally with petrologic type  $3 > 4 > 5 \geq 6$ . When normalized to Ni, less-depleted elements vary with chondritic groups in UOC. Examination of statistically-significant interelement relationships among various ordinary chondrite populations involving 34 elements reveals patterns distinct from those of other chondritic groups. These patterns reflect nebular metal-silicate fractionation which preceded or accompanied thermal fractionation. Metamorphic effects are not pronounced among volatile/mobile elements in UOC and, although condensation models seem generally more satisfactory, none successfully accounts for all observations.

Thus it appears that primitive chondrites originated in several ways: carbonaceous chondrites by the mixing of two components differing in thermal history; enstatite chondrites by a nebular thermal fractionation accompanied by a sulfide-plus-metal separation from silicate and followed by metamorphism in a parent body; and UOC by a nebular thermal fractionation accompanied by metal separation from silicate. It seems that there is no unique process by which primitive solar material condensed to form early objects.

In paper (6) cited in Part I we report our determinations of Ag, Bi, Cd, Co, Cs, Ga, In, Se, Te, Tl and Zn in the 6 known ureilites. All 11 elements are depleted below cosmic, i.e. Cl, levels and their characteristic abundance pattern differs substantially from those of the chondritic groups. These and other data can be treated under a variety of assumptions and all treatments lead to the same conclusion: ureilites do not derive from a simple mixture

of volatile-rich chondrites with achondritic material. We suggest, rather, that they derive from a mixture of cosmochemically-fractionated achondritic material (similar but not identical to Chassigny) and a "distillate" of thermally mobile elements which was then shocked during breakup of the parent object.

### III. Other Studies


We are continuing our studies of shock effects in meteorites at a low-priority level. We are now preparing a manuscript describing our shock- and cosmogenic gas-study (with Professor G. Herzog of Rutgers University) of samples from known geographic locations around the Odessa crater.

### IV. Administrative Information:

The following individuals are co-workers at Purdue currently doing research on this grant:

- A. Dr. C. M. Binz, Postdoctoral Research Associate
- B. Dr. M. Ikramuddin, Postdoctoral Research Associate
- C. Mr. S. Matza, Graduate Assistant in Research
- D. Mr. J. Carvalho, Graduate Research Assistant

The NASA Technical Officer for this grant is Dr. D. Stuart-Alexander, NASA Headquarters, Lunar Programs Office, Code SM, Washington, D. C., 20546.

  
Michael E. Lipschutz

cc: Dr. Bryson, NASA

Dr. Stuart-Alexander, NASA

Mr. Baitinger, Purdue

Dr. Waling, Purdue